


Docket No. 2000-096DIV5
PATENTIN THE CLAIMS

Prior to issuance, please make the following amendments to correct typographical errors in the claims:

1. (Currently amended) A metal-ligand complex characterized by the following formula:



wherein R^1 is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl and combinations thereof;


T is a bridging group selected from the group consisting of $-\text{CR}^2\text{R}^3-$ and $-\text{SiR}^2\text{R}^3-$ with R^2 and R^3 independently selected from the group from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, silyl, boryl, phosphino, amino, thio, seleno, halide, nitro, and combinations thereof;

J''' is selected from the group consisting of heteroaryl and substituted heteroaryl;

each L is independently selected from the group consisting of halide, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, hydroxy, boryl, silyl, amino, amine, hydrido, allyl, diene, seleno, phosphino, phosphine, carboxylates, thio, 1,3-dionates, oxalates, carbonates, nitrates, sulphates, ethers, thioethers and combinations thereof or optionally two or more L groups are joined into a ring structure; n is 1, 2, 3, 4, 5, or 6; and x is 1 or 2.


Docket No. 2000-096DIV5
PATENT

2. (Original) The metal complex of claim 1 having the formula:



wherein each of R^4 , R^5 , R^6 and R^7 is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, silyl, boryl, phosphino, amino, thio, seleno, halide, nitro, and combinations thereof; and optionally, any combination of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 or R^7 may be joined together in a ring structure.

3. (Original) The metal complex of claim 2, wherein said complex is characterized by the formula:




wherein Q^1 , Q^2 , Q^3 , Q^4 and Q^5 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, silyl, boryl, phosphino, amino, thio, seleno, nitro,

Docket No. 2000-096DIV5
PATENT

and combinations thereof; optionally any 2 of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 are joined together in a ring structure; and $x = 1$.


4. (Original) The metal complex of claim 2, wherein said complex is characterized by the formula:



such that T is $-CR^2R^3-$ and wherein R^{10} , R^{11} , R^{12} and R^{13} are each independently selected from the group consisting of hydrogen, halide, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, silyl, boryl, phosphino, amino, thio, seleno, nitro, and combinations thereof; optionally, two or more R^{10} , R^{11} , R^{12} and R^{13} groups may be joined to form a fused ring system having from 3-50 non-hydrogen atoms; and


R^{14} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, silyl, boryl, phosphino, amino, thio, seleno, halide, nitro, and combinations thereof; and $x = 1$.

5. (Currently amended) The metal complex of claim 4, wherein said complex is characterized by the general formula:

Docket No. 2000-096DIV5
PATENT

hereinwherein Q^1 , Q^2 , Q^3 , Q^4 and Q^5 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, silyl, boryl, phosphino, amino, thio, seleno, nitro, and combinations thereof; optionally any 2 of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 are joined together in a ring structure.

6. (Currently amended) A metal complex characterized by the formula:



where M is zirconium or hafnium;

wherein R^1 is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl and combinations thereof;

T is a bridging group selected from the group consisting of $-CR^2R^3-$ and $-SiR^2R^3-$ with R^2 and R^3 being independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl,


Docket No. 2000-096DIV5
PATENT

heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxyl, aryloxyl, silyl, boryl, phosphino, amino, thio, seleno, halide, nitro, and combinations thereof;

J''' being selected from the group of substituted heteroaryls with 2 atoms bonded to the metal M, at least one of those 2 atoms being a heteroatom, and with one atom of J''' is bonded to M via a dative bond, the other through a covalent bond; and

L¹ and L² are independently selected from the group consisting of halide, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, hydroxy, boryl, silyl, amino, amine, hydrido, allyl, diene, seleno, phosphino, phosphine, carboxylates, thio, 1,3-dionates, oxalates, carbonates, nitrates, sulphates, ethers, thioethers and combinations thereof or optionally the two L groups are joined into a ring structure.

7. (Original) The complex of claim 6, wherein said complex is characterized by the formula:




wherein each of R⁴, R⁵ and R⁶ is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxyl, aryloxyl, silyl, boryl, phosphino, amino, thio, seleno, halide, nitro, and combinations thereof; and optionally, any combination of R¹, R², R³, R⁴, R⁵, or R⁶ may be joined together in a ring structure; and

E'' is either carbon or nitrogen and is part of a cyclic aryl, substituted aryl, heteroaryl, or substituted heteroaryl group.


Docket No. 2000-096DIV5
PATENT

8. (Original) The metal complex of claim 7, wherein said complex is characterized by the formula:



wherein R^{10} , R^{11} , R^{12} and R^{13} are each independently selected from the group consisting of hydrogen, halide, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxy, aryloxy, silyl, boryl, phosphino, amino, thio, seleno, nitro, and combinations thereof; optionally, two or more R^{10} , R^{11} , R^{12} and R^{13} groups may be joined to form a fused ring system having from 3-50 non-hydrogen atoms.

9. (Original) The metal complex of claim 8, wherein said complex is characterized by the formula:




wherein Q^1 , Q^2 , Q^3 , Q^4 and Q^5 are independently selected from the group consisting of

Docket No. 2000-096DIV5
PATENT

hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxyl, aryloxyl, silyl, boryl, phosphino, amino, thio, seleno, nitro, and combinations thereof; or optionally, two of Q^2 , Q^3 and Q^4 are joined together in a ring structure.

10. (Currently amended) The complex of either of claims 1, 2, 4, 6, 7 or 8, wherein R^1 is characterized by the general formula:




wherein E is either carbon or nitrogen,

Q^1 and Q^5 are substituents on the R^1 ring at a position ortho to E, with at least one of Q^1 or Q^5 being bulky, wherein Q^1 and Q^5 are independently selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl and silyl, but provided that Q^1 and Q^5 are not both methyl; and

Q''_q represents additional possible substituents on the ring, with q being 1, 2, 3, 4 or 5 and Q'' being selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxyl, aryloxyl, silyl, boryl, phosphino, amino, thio, seleno, halide, nitro, and combinations thereof.

11. (Original) The complex of claim 10, wherein R^1 is characterized by the general formula












Docket No. 2000-096DIV5
PATENT

wherein Q^2 and Q^3 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heteroalkyl, substituted heteroalkyl, heterocycloalkyl, substituted heterocycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, alkoxyl, aryloxy, silyl, boryl, phosphino, amino, thio, seleno, nitro, and combinations thereof; or optionally, Q^2 and Q^3 are joined together in a ring structure.

12. (Original) The complex of either of claims 6, 7, 8 or 9, wherein M is hafnium.

13. (Original) The complex of either of claims 6, 7, 8 or 9, wherein L^1 and L^2 are the same and selected from the group consisting of methyl and dimethylamino.

14. (Original) The complex of claim 13, wherein said complex is selected from the group consisting of:



Docket No. 2000-096DIV5
PATENT

15. (Original) A catalyst comprising the complex of either of claims 1, 2, 3, 4, 5, 6, 7, 8 or 9, combined with an activator, combination of activators or activating technique.

16. (Currently amended) The catalyst of claim 15, wherein said catalyst comprises an activator that is an ionic activator.

17. (Original) The catalyst of claim 16, additionally comprising at least one of a group 13 reagent, a divalent metal reagent or an alkali metal reagent.

NO FURTHER AMENDMENTS ON THIS PAGE